

- Golyshev, S. I., and Padalko, N. L. (1979) A theoretical study of oxygen-isotope distribution in oxides. *Geochem. Int.* **16**, 157–60.
- Harzer, D. (1970) Geochemical studies on hydrothermal minerals from deposits in the German Democratic Republic with the isotopes oxygen-18 and carbon-13 (in German). *Freiberg, Forschungsh. C*, **247**, 132 p.
- Hattori, K., and Halas, S. (1982) Calculation of oxygen isotope fractionation between uranium dioxide, uranium trioxide and water. *Geochim. Cosmochim. Acta*, **46**, 1863–8.
- Jackson, N. J., and Rankin, A. H. (1976) Fluid inclusion studies at St Michael's Mount. *Proc. Ussher Soc.* **3**, 430–4.
- Kelly, W. C., and Rye, R. O. (1979) Geologic, fluid inclusion, and stable isotope studies of the tin-tungsten deposits of Panasqueira, Portugal. *Econ. Geol.* **74**, 1721–822.
- Kieffer, S. W. (1982) Thermodynamics and lattice vibrations of minerals: 5. Applications to phase equilibria, isotopic fractionation, and high pressure thermodynamic properties. *Rev. Geophys. Space Phys.* **20**, 827–49.
- Matthews, A. (1973) *Oxygen isotope geology: Experimental calibration of geothermometers with related experimental investigations*. Unpubl. PhD thesis, Univ. of Manchester.
- Matsuhisa, Y., Goldsmith, J. R., and Clayton, R. N. (1979) Oxygen isotope fractionation in the system quartz-albite-anorthite-water. *Geochim. Cosmochim. Acta*, **43**, 1131–40.
- Moore, F., and Moore, D. J. (1979) Fluid inclusion study of mineralization at St Michael's Mount, Cornwall, England. *Trans. Instn Min. Metall. (B)*, **88**, 57–60.
- Mu Zhigo *et al.* (1982) Oxygen, hydrogen and carbon isotope studies of Piaotang and Xihuashan quartz vein-type tungsten deposits, Jiangxi Province (in) Tungsten Geology symposium (Jiangxi, China, Oct. 1981), ESCAP/RMRDC (Bandung) and Geol. Publ. House (Beijing), 385–401.
- O'Neil, J. R. (1986) Theoretical and experimental aspects of isotopic fractionation. In *Stable isotopes* (Valley, J. W., Taylor, H. P., and O'Neil, J. R., eds.) *Reviews in Mineralogy*, **16**, Mineral. Soc. America, 1–40.
- Patterson, D. J., Ohmoto, H., and Solomon, M. (1981) Geologic setting and genesis of cassiterite-sulfide mineralization at Renison Bell, Western Tasmania. *Econ. Geol.* **76**, 393–408.
- Sun, S-S., and Eadington, P. J. (1987) Oxygen isotope evidence for the mixing of magmatic and meteoric waters during tin mineralization in the Mole Granite, New South Wales, Australia. *Ibid.* **83**, 43–52.
- Sushchevskaya, T. M., Ustinov, V. I., Nekrasov, I. A., Gavrilov, Y. Y., and Grinenko, V. A. (1986) The oxygen-isotope fractionation factor in cassiterite synthesis. *Geochem. Int.* **23**, 57–60.

[Manuscript received 21 April 1988;  
revised 20 September 1988]

© Copyright the Mineralogical Society

KEYWORDS: oxygen isotopes, cassiterite, water, fractionation.

Department of Geology,  
RHB New College,  
Egham,  
Surrey, TW20 0EX

DAVID H. M. ALDERTON

MINERALOGICAL MAGAZINE, JUNE 1989, VOL. 53, PP. 376–379

## Högbomite from the Aldan Shield, Eastern Siberia, USSR

HÖGBOMITE, a complex oxide of Al, Fe, Mg, and Ti, is an important constituent of some iron ores and emery deposits as well as an infrequent accessory in aluminous high-grade rocks (e.g. Grew *et al.*, 1987). The recent increase in reports of new localities (e.g. Rammlmair *et al.*, 1988) suggests that högbomite may be more widespread than is generally perceived. We report here högbomite from the Aldan Shield, Eastern Siberia. This högbomite is remarkable for the wide variation in composition measured in a single thin section. Our report is only the second from the USSR of högbomite for which chemical data are given. Reports of högbomite other than Moleva and Myasnikova's (1952) well-documented descrip-

tion of högbomite from the Urals are based only on optical properties (Bobrovnik, 1955; Sinita, 1957; Sudovikov *et al.*, 1962); the third citation includes mentions of högbomite in pelitic gneisses from unspecified localities in the Aldan Shield. In the present paper we present details of the paragenesis and chemistry of högbomite and several associated minerals; the reader is referred to Drugova *et al.* (in press) for a general account of the högbomite-bearing rock and its geological and geochemical significance.

Högbomite is found in a biotite-plagioclase-corundum-spinel schist cropping out 2 km east of the Aldan River at a point 1 km north of its tributary Ayyannaakh Creek (approximately 120 km

southwest of the city of Aldan). The h ogbomite-bearing rock is part of a metasedimentary unit consisting largely of quartzite with layers of pelitic gneisses and schists; associated rocks include orthopyroxene-bearing plagiogneiss. These rocks were metamorphosed in the granulite facies, subsequently migmatized during two events in the amphibolite facies, and intruded by microcline granite (Drugova *et al.*, in press).

The h ogbomite-bearing rock is a biotite-plagioclase schist containing well-aligned cylindrical nodules of corundum, magnetite, and spinel up to 10 cm long and 6 cm across as well as disseminated grains of corundum, magnetite, rutile(?) and h ogbomite. The oxide nodules constitute up to 30 or 40% of the rock and are found over an area of 20 or 30 m<sup>2</sup> in outcrop. The matrix to the nodules is a medium-grained rock (mostly 0.5–3 mm) consisting of dark biotite, plagioclase, subordinate magnetite, rare clinozoisite, and minor secondary muscovite. The plagioclase (An 40–46 in the section analysed with the microprobe, Drugova *et al.*, in press) is locally zoned. The nodules generally have a core of corundum up to 12 mm across, surrounded by a mantle of spinel, magnetite, and corundum. Adjacent to the corundum core is an inner rim of magnetite. Spinel is commonly dominant in the remainder of the mantle; magnetite, rarely in association with ilmenite, occurs as isolated grains. The spinel also contains abundant blebs of exsolved magnetite in distinct planar arrays. Towards the margin of the mantle are biotite and patches of corundum riddled with magnetite. This corundum is in places dichroic in blue. In the analysed section, the oxide nodule has a corona of fine-grained pale biotite. Discrete biotite flakes near this corona are in places pale brown or green. Other minerals in the nodule-bearing rocks are apatite, zircon, allanite, monazite, pyrrhotite, and rutile (Drugova *et al.*, in press); of these, only zircon was found in the analysed section.

H ogbomite is found in both the matrix and in the nodules; its distribution is irregular. Grains are mostly 0.05 to 0.2 mm across, locally up to 0.35 mm, and range from nearly equant to planar. In the matrix, h ogbomite is closely associated with magnetite and forms aggregates contiguous to irregular magnetite grains; in some cases, h ogbomite appears to have replaced magnetite; in others, textural relations are ambiguous. Near some h ogbomite, magnetite appears to be replaced by a fine-grained mineral resembling rutile (leucocene?). In the nodules, h ogbomite occurs mostly in and near spinel and locally with magnetite or corundum and, in places, touches biotite. H ogbomite is moderate brown (nodule) to dark

brown (matrix) and is dichroic.

One grain of the nodule h ogbomite and three grains (in two aggregates) of the matrix h ogbomite, together with spinel (in two spots) and associated minerals, were analysed in a single thin section within a 1 × 1.5 cm area with a Cameca 'Camebax-micro' electron microprobe at the Institute of Geology, Yakutsk (selected analyses listed Table 1). The nodule and matrix h ogbomite differ most in Ti, total Fe, and Mg. Matrix h ogbomites also vary in composition from grain to grain and within a grain (e.g. TiO<sub>2</sub>, 4.6–5.6%; ZnO, 0.8–1.8%), and the nodule h ogbomite grain also appears to be zoned in Ti. In formulae recalculated assuming 22 cations, 30 oxygens and 2 hydroxyls (Gatehouse and Grey, 1982; Grew *et al.*, 1987), most of the Fe in the matrix h ogbomite is Fe<sup>2+</sup> and that in the nodule h ogbomite, Fe<sup>3+</sup>, and the two compositions can be related by the substitutions  $Ti + Fe^{2+} = 2Fe^{3+}$  ( $TiFe_{-1}$ , ilmenite-hematite type) and  $Mg = Fe$  ( $MgFe_{-1}$ ); Al contents in the formulae are nearly identical. However, these formulae were calculated assuming the 8H polytype, apparently a rare polytype (Beukes *et al.*, 1986), and thus may not be valid for either or both of the Aldan h ogbomites. Indeed, the excessively high analytical totals calculated for the nodule h ogbomites (e.g. 102.4%, Table 1) indicate that this calculation results in an excessive estimate of Fe<sup>3+</sup> content. On the other hand, the Zakrzewski (1977) formula for recasting h ogbomite analyses yields Fe<sup>3+</sup>/Total Fe ratios of 0.07 and 0.08 for the matrix h ogbomite and <0 and 0.05 for the nodule h ogbomite. These values seem too low, because the corresponding ratios in spinel are calculated to be 0.10 to 0.13. In the absence of detailed crystallographic studies of different h ogbomite polytypes, it is not possible to reliably recast analyses and estimate Fe<sup>3+</sup>/Fe<sup>2+</sup> ratios, nor is it possible to determine water contents. We have calculated a value of H<sub>2</sub>O using the Gatehouse and Grey (1982) formula in order to call attention to this potentially critical component of h ogbomite.

Magnetite (5 analyses) in the matrix and nodules contain 0.2–0.3% Cr<sub>2</sub>O<sub>3</sub> and 0–0.2% TiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub>, MnO, MgO, and ZnO (for example, Table 1), with the exception of one spot with 3.1% Al<sub>2</sub>O<sub>3</sub> in the matrix. The averaged spinel analyses in Table 1 are representative of the 4 analyses in Cr<sub>2</sub>O<sub>3</sub> contents (0.08–0.12%) and  $X(Fe)T(0.35–0.36)$ , while ZnO in the unlisted analyses reaches 1.68%. Corundum (4 analyses) in the spinel mantle contains 0.37–0.99% Fe<sub>2</sub>O<sub>3</sub> and 0.18–0.24% Cr<sub>2</sub>O<sub>3</sub>. The dark biotites of the matrix (4 flakes analysed; average of 2 in Table 1) are titanian (2.8–3.1% TiO<sub>2</sub>), and relatively

Table 1. Composition of Minerals in Sample 0206B, Ayyannaakh Stream, Aldan Shield, USSR

	Biotite			Högbomite		Spinel	Magnetite		Corundum	
	Matrix	Corona	Corona	Matrix	Nodule	Nodule	Nodule	Matrix	Nodule	
Number of Analyses	5	1	1	1	1	2	2	1	1	1
SiO <sub>2</sub>	35.82	37.16	38.48	0.0	0.05	0.01	0.0	0.0	0.0	0.0
TiO <sub>2</sub>	3.12	0.38	0.24	5.30	5.55	2.95	0.0	0.0	0.0	0.0
Al <sub>2</sub> O <sub>3</sub>	17.57	21.52	19.10	61.20	60.74	64.67	65.40	0.21	0.15	99.04
Cr <sub>2</sub> O <sub>3</sub>	0.06	0.07	0.06	0.25	0.20	0.17	0.10	0.18	0.29	0.19
FeO <sup>1</sup>	13.54	7.22	7.43	24.26	25.30	17.10	17.29	87.88	87.74	0.99
MnO	0.11	0.12	0.08	0.25	0.34	0.27	0.39	0.0	0.03	0.01
MgO	13.65	19.43	20.48	5.97	5.47	13.14	17.57	0.0	0.0	0.01
ZnO	—	—	—	1.35	0.83	0.94	0.63	0.15	0.03	—
CaO	0.0	0.03	0.02	0.0	0.01	0.0	0.0	0.0	0.0	0.02
Na <sub>2</sub> O	0.15	0.16	0.13	0.07	0.02	0.02	0.03	0.08	0.0	0.0
K <sub>2</sub> O	10.36	10.30	10.04	0.0	0.05	0.0	0.0	0.02	0.0	0.0
H <sub>2</sub> O <sup>2</sup>	3.98	4.22	4.21	1.45	1.45	1.55	—	—	—	—
Total	98.36	100.61	100.27	100.10 <sup>3</sup>	100.01 <sup>4</sup>	100.82 <sup>5</sup>	101.41 <sup>6</sup>	88.52 <sup>7</sup>	88.24 <sup>8</sup>	100.26
Formulae										
Oxygens	22	22	22	31	31	31	4	4	4	3
Si	5.398	5.278	5.478	0.0	—	—	0.0	0.0	0.0	0.0
Al <sup>IV</sup>	2.602	2.722	2.522	—	—	—	—	—	—	—
Al	—	—	—	14.871	14.839	14.790	1.950	0.010	0.007	1.985
Al <sup>VI</sup>	0.518	0.880	0.683	—	—	—	—	—	—	—
Cr	0.007	0.008	0.007	0.041	0.033	0.026	0.002	0.006	0.009	0.003
Ti	0.354	0.041	0.026	0.822	0.865	0.430	0.0	0.0	0.0	0.0
Fe <sup>2+</sup>	1.706	0.858	0.885	2.738	2.988	0.452	0.317	0.996	0.998	—
Fe <sup>3+</sup>	—	—	—	1.445	1.398	2.323	0.048	1.984	1.983	0.013
Mn	0.014	0.014	0.010	0.044	0.060	0.044	0.008	0.000	0.001	—
Mg	3.067	4.114	4.346	1.835	1.690	3.800	0.663	0.000	0.000	—
Total	5.666	5.915	5.957	—	—	—	—	—	—	—
Ca	0.0	0.005	0.004	—	—	—	—	—	—	—
Na	0.045	0.044	0.036	—	—	—	—	—	0.0	—
K	1.991	1.866	1.823	—	—	—	—	—	0.0	—
Total	2.036	1.915	1.863	—	—	—	—	—	—	0.0
Zn	—	—	—	0.206	0.127	0.134	0.012	0.004	0.001	—
Total	15.702	15.830	15.820	22.002	22.000	21.999	3.0	3.000	2.999	2.001
X(Fe)T <sup>9</sup>	0.36	0.17	0.17	0.70	0.72	0.42	0.36	—	—	—

## Notes.

Dash - Not analysed or not calculated.

- All Fe as FeO, except for corundum, for which all Fe as Fe<sub>2</sub>O<sub>3</sub>. In formulae, Fe<sup>2+</sup> and Fe<sup>3+</sup> calculated from stoichiometry.
- Calculated assuming ideal hydroxyl contents for högbomite (2(OH) and 22 cations, Gatehouse and Grey, 1982) and for biotite.
- 3-8. Calculated weight %: 3- FeO=15.88, Fe<sub>2</sub>O<sub>3</sub>=9.31; Total=101.03, 4- FeO=17.24, Fe<sub>2</sub>O<sub>3</sub>=8.96, Total=100.91; 5- FeO=2.79, Fe<sub>2</sub>O<sub>3</sub>=15.91, Total=102.42; 6- FeO=15.00, Fe<sub>2</sub>O<sub>3</sub>=2.54, Total=101.66; 7- FeO = 29.36, Fe<sub>2</sub>O<sub>3</sub> = 65.04, Total = 95.04; 8- FeO = 29.37, Fe<sub>2</sub>O<sub>3</sub> = 64.87, Total = 94.74.
- X(Fe)T = total Fe/(Mg + Total Fe)

Microprobe Operating Conditions: 20 kV, 40 nA.

iron rich ( $X_{\text{Fe}} = 0.35\text{--}0.37$ ), while the fine-grained biotites in the corona (7 analyses) are almost free of Ti (0–0.4%  $\text{TiO}_2$ ), aluminous (19–23%  $\text{Al}_2\text{O}_3$ ) and magnesian ( $X_{\text{Fe}} = 0.15\text{--}0.17$ ).

The present mineralogy and chemistry of the h ogbomite-bearing rocks is due to a complex metamorphic history to which Drugova *et al.* (in press) assigned 3 stages. The  $X_{\text{Fe}}$ 's of the matrix biotite, which is interpreted to have formed during the granulite-facies event, and spinel are approximately equal, that is  $K_D = [\text{Fe}/\text{Mg}(\text{Sp}1)]/[\text{Fe}/\text{Mg}(\text{Bt})] = 0.91\text{--}1.00$ , in marked contrast to equilibrium spinel–biotite pairs, for which  $K_D \geq 3.7$  (Lal *et al.*, 1978; Waters and Moore, 1985). Consequently, the analysed spinel probably did not equilibrate with the matrix biotite, whose rather constant composition from grain to grain is presumed to be little changed since original crystallization in the granulite facies. The spinel–magnetite–corundum intergrowths in the mantles of the nodules could have resulted from exsolution, breakdown, and possibly oxidation of a homogeneous hercynite-rich member of the spinel group that had equilibrated with the matrix biotite. H ogbomite in the nodule probably formed during this breakdown, drawing on Ti and Zn in the original homogeneous spinel. By analogy with other h ogbomite occurrences (e.g. Grew *et al.*, 1987), appearance of h ogbomite suggests interaction of the secondary spinel, magnetite, and corundum with hydrous fluids. Alternatively Ti for h ogbomite could have originated from the matrix biotite, which released Ti, its remaining constituents forming the corona biotite. The  $K_D$  (Mg–Fe) of 2.5–3.1 for spinel and corona biotite is closer to the presumably equilibrium values cited above. Thus h ogbomite formation in the nodules, breakdown of a hercynite-rich phase into spinel, magnetite, and corundum, and crystallization of corona biotite could be coeval and closely related to either intrusion of microcline granite or retrogression in the amphibolite facies.

The matrix h ogbomite is presumed to have formed coevally with the nodule h ogbomite and could have formed from magnetite, but there is no obvious source of Al. Alternatively the present magnetite–h ogbomite  $\pm$  rutile (?) aggregates

KEYWORDS: h ogbomite, Aldan Shield, Siberia, USSR.

could have resulted from the recrystallization (and oxidation?) of a pre-existing Fe–Mg–Ti–Al  $\pm$  Zn spinel phase (+ ilmenite?). In any case, given its different composition, the matrix h ogbomite must have formed from a different mix of oxide phases from those in the nodule, thereby implying that compositional factors, including oxygen fugacity, rather than pressure and temperature, are the main controls on h ogbomite chemistry.

This research was carried out with the support of the interacademy exchange program between the USSR and USA within the framework of IGCP Project 235 'Metamorphism and Geodynamics'.

### References

- Beukes, G. J., van Zyl, V. C., Schoch, A. E., De Bruijn, H., van Aswegen, G., and Strydom, D. (1986) *Neues Jahrb. Mineral. Abh.* **155**, 53–66.
- Bobrovnik, D. P. (1955) In *Voprosy mineralogii osadochnykh obrazovaniy*, Kniga 2-ya [Problems of mineralogy of sedimentary formations, Book 2], L'vovskiy Gosud. Univ. im. Ivana Franko, 215–16.
- Drugova, G. M., Grew, E. S., and Leskova, N. V. *Zap. Vses. mineral. Obshch.* (in press).
- Gatehouse, B. M., and Grey, I.E. (1982) *Am. Mineral.* **67**, 373–80.
- Grew, E. S., Abraham, K., and Medenbach, O. (1987) *Contrib. Mineral. Petrol.* **95**, 21–31.
- Lal, R. K., Ackermann, D., Seifert, F., and Haldar, S. K. (1978) *Ibid.* **67**, 169–87.
- Moleva, V. A., and Myasnikov, V. S. (1952) *Dokl. Akad. Nauk SSSR*, **53**, 733–36.
- Rammlmair, D., Mogessie, A., Purtscheller, F., and Tessadri, R. (1988) *Am. Mineral.* **73**, 651–6.
- Sinita, S. (1957) *Mineral. Sbornik L'vovsk. Geol. Obshch.* **11**, 170–86.
- Sudovikov, N. G., Drugova, G. M., Krylova, M. D., and Mikhaylov, D. A. (1962) In *Petrografia Vostochnoy Sibiri* [Petrography of Eastern Siberia], vol. 2, Moscow, Acad. Sci. Publishing House, 320–446.
- Waters, D. J., and Moore, J. M. (1985) *Contrib. Mineral. Petrol.* **91**, 369–82.
- Zakrzewski, M. A. (1977) *Neues Jahrb. Mineral. Mh.*, 373–80.

[Manuscript received 29 June 1988;  
revised 10 October 1988]

  Copyright the Mineralogical Society

Department of Geological Sciences, University of Maine, Orono, ME 04469, USA EDWARD S. GREW

Institute of Geology and Geochronology of the Precambrian, Academy of Sciences of the USSR, Leningrad, USSR 199164 GALINA M. DRUGOVA

Institute of Geology, Yakutsk Filial, Siberian Division, Academy of Sciences of the USSR, Yakutsk, USSR N. V. LESKOVA