

M. NAKATA AND H. HONMA

*Department of Astronomy and Earth Science,
Tokyo Gakuji University, Koganei-shi,
Tokyo 184, Japan*

J. I. CHUNG

*Department of Geology, College of Natural Science,
Chonbug University, Chollabuk-do, Korea*

K. SAKURAI

*Sakurai Museum, Hirakawa-cho, Chiyoda-ku,
Tokyo 105, Japan*

[Manuscript received 23 March 1988;
revised 10 June 1988]

© Copyright the Mineralogical Society

Wickmanite from Whealcock Zawn, Botallack, Cornwall

THE rare mineral wickmanite, $Mn^{2+}Sn^{4+}(OH)_6$, was first described from Långban, Sweden (Moore and Smith, 1967), where it occurs as brown to honey yellow octahedra up to 2 mm in size in pockets in magnetite ore and in a brecciated jacobsonite-richterite-manganophyllite skarn. It was later found in a nepheline-syenite pegmatite at Tvedalen, Norway (Amli and Griffin, 1972), associated with leadhillite and hydrocerussite; at Pitkaranta, Karelia, USSR (Nefodov *et al.*, 1977), in a hydrothermally mineralised skarn; and at Llallagua, Bolivia (Kampf, 1982), where it occurs in pockets in stannite. It has now been found at Whealcock Zawn, Botallack, Cornwall, the first recorded occurrence in the UK, having been confirmed at the British Museum (Natural History) by means of X-ray powder diffraction (film no. 5452F).

At Whealcock Zawn the wickmanite occurs as orange-yellow octahedra, up to 1 mm on edge, showing parallel growth. The wickmanite is found in small cavities in a large axinite pod in a discordant calc-silicate body (Alderton and Jackson, 1978) exposed on the south side of Whealcock Zawn (SW 363 340) which is within the metamorphic aureole of the Lands End granite.

The associated minerals include abundant massive and more rarely euhedral grossular, coarsely bladed dark green pargasite, white prisms of apatite to 50 mm, small white rosettes of titanite, rare pinkish crystals of orthoclase (variety adularia), minor chalcocopyrite which is locally altered to botallackite (Barstow, pers. comm.), and an unidentified cobalt-bearing mineral.

It is interesting to note that the locality has also produced crystals of the rare tin silicate stokesite (Couper and Barstow, 1977; Couper and Clark, 1977) and may indeed be the type locality for that mineral.

Acknowledgements. Thanks are due to J. G. Francis of the Department of Mineralogy, British Museum

(Natural History) for the X-ray powder diffraction determination.

REFERENCES

- Alderton, D. H. M., and Jackson, N. J. (1978) *Mineral. Mag.* **42**, 427–34.
 Amli, R., and Griffin, W. L. (1972) *Nor. Geol. Tidsskr.* **52**, 193–6.
 Couper, A. G., and Barstow, R. W. (1977) *Mineral. Rec.* **8**, 294–7.
 — and Clark, A. M. (1977) *Mineral. Mag.* **41**, 411–4.
 Kampf, A. R. (1982) *Mineral. Rec.* **13**, 235–9.
 Moore, P. B., and Smith, J. V. (1967) *Arkiv. Min. Geol.* **4**, 395–9.
 Nefodov, E. I., Griffin, W. L., and Kristiansen, R. (1977) *Can. Mineral.* **15**, 437–45.

KEYWORDS: wickmanite, stokesite, Botallack, Cornwall.

NEIL HUBBARD

122 Cordery Road, Evington, Leicester, LE5 6DF, UK

[Manuscript received 12 August 1988]

© Copyright the Mineralogical Society

Cinnabar from the northern Pennines, England

THE first record of British cinnabar is that of Braithwaite *et al.* (1963) who described the mineral from Rutland Cavern, Matlock Bath, Derbyshire. A. W. G. Kingsbury subsequently identified cinnabar from several places on Grassington Moor and Greenhow Hill where, in a few places, the other mercury minerals metacinnabarite, calomel, and native mercury were also found (Dunham and Wilson, 1985). Traces of cinnabar have been noted in pan concentrates of stream sediments from two tributaries of the River Clyde near Abington, Lanarkshire (Dawson *et al.*, 1979) and from streams near the Glendinning antimony deposit, Langholm, Dumfriesshire (Gallagher *et al.*, 1983). Cinnabar is described here as a rare supergene mineral from six localities in the Northern Pennines.

The minerals were identified by X-ray diffraction in the Department of Geology, University of Sheffield.

Coldstones Quarry, Pateley Bridge, North Yorkshire [SE 123 641]. The southern face of this quarry exposes the Garnet Vein (Dunham and Wilson, 1985, p. 209), which here is up to 2 m wide and composed mainly of fluorite, baryte, and calcite with small amounts of galena. Supergene minerals include hemimorphite, smithsonite and traces of aurichalcite and rosasite (D. Green, pers. comm.). In places, irregular cavities lined with smithsonite, hemimorphite, and calcite appear to represent original pockets of primary minerals, probably mainly sphalerite. Cinnabar occurs within these cavities as bright orange-red earthy coatings on the other