

pressure-temperature-time paths, 799 pp.
 Monograph M.S.A, Washington D.C.
 Vallance, T.G. (1967) *J. Petrol.*, 8, 84-96.

[Manuscript received 18 January 1995;
 revised 29 March 1995]

© Copyright the Mineralogical Society

KEYWORDS: subsilicic sodium gedrite, quartz keratophyric protolith, Nordmark, Sweden.

MINERALOGICAL MAGAZINE, APRIL, VOL. 60, PP 387-388

Clinobisvanite, eulytite, and namibite from the Pala pegmatite district, San Diego Co., California, USA

Eugene E. Foord

MS 905, U.S. Geological Survey,
 Box 25046 Denver Federal Center,
 Denver, CO 80225,
 USA

THE 100 Ma complex LCT-type composite pegmatite-aplite dykes, intruded into various units of the Southern California Batholith, are known to contain bismuth minerals. Jahns and Wright (1951) reported the following primary and secondary bismuth minerals from the quartz-rich cores of a number of dykes in the Pala district, San Diego Co., California: native bismuth, bismuthinite, bismite, bismutite, and beyerite. The Little Three pegmatite-aplite dyke in the Ramona district, contains minor amounts of native bismuth, beyerite, bismite, bismutite, and pucherite (Shigley *et al.*, 1986; Foord *et al.*, 1989). The Himalaya dyke system in the Mesa Grande district contains bismuth, bismutite, pucherite and beyerite. All of these minerals are sparse but characteristic of highly evolved granitic pegmatites.

Additional mineralogical studies in the Pala district indicate that at least three additional bismuth minerals not reported by Jahns and Wright (1951) or Jahns (1979) are present. The occurrence of namibite from Bohemia was recently reported by Mrázek *et al.* (1994). Namibite was reported in 1991 from the Pala district by Foord *et al.* (1991).

All minerals were identified by X-ray diffraction, SEM-EDS analyses and by electron-microprobe analyses. Fine-grained emerald-green namibite intergrown with fine-grained, tan clinobisvanite and beyerite occurs in small pod-like masses in quartz-rich 'pocket-pegmatite' in the White Queen mine on Hiriart Mountain. A total of several tens of grams of

material was recovered. Electron-microprobe analyses of the clinobisvanite and namibite are given in Table 1.

An 0.5 kg mass of red-orange crystalline (individual crystals to several mm long) clinobisvanite and cream-white to tan radiating hemispheres (to several mm) of eulytite, $\text{Bi}_4(\text{SiO}_4)_3$, and associated albite occurring in quartz-rich core-zone pegmatite was found in the Elizabeth "R" mine on Chief Mountain. While eulytite is known to be cubic, and occurring in hextetrahedral crystals, the radiating and semi-globular aggregate habit agrees perfectly with that of material from both Schneeberg and

TABLE 1. Electron-microprobe analyses for clinobisvanite and namibite from the White Queen mine

	Clinobisvanite (ave. of five analyses)	Namibite (ave. of five analyses)
V ₂ O ₅	27.0	14.1
CuO	—	12.5
Bi ₂ O ₃	72.0	71.9
CaO	0.9**	0.1
H ₂ O	—	1.4*
Total	99.9	100.0

* —by difference. ** — minor beyerite contaminant.

Johanngeorgenstadt, Saxony (Fron del, 1943). Small amounts of namibite with associated chrysocolla and pseudomalachite also were found in quartz-rich pegmatite in the Stewart mine on Tourmaline Queen Mountain.

Three polymorphs of BiVO_4 are known: clinobisvanite (monoclinic), pucherite (orthorhombic), and dreyerite (tetragonal). Only the first two have been found thus far in San Diego Co.

Empirical formulae for the two minerals are as follows: (using 4 oxygen atoms for clinobisvanite and 7 oxygen atoms for namibite) $\text{Bi}_{1.01}\text{V}_{0.97}\text{O}_4$ and $\text{Cu}_{1.01}(\text{BiO})_{1.99}\text{V}_{1.00}\text{O}_4(\text{OH})_{1.00}$ respectively. Electron-microprobe analyses were not made of the eulytite from the Elizabeth R mine. The revised formula for namibite proposed by Mrázek *et al.* (1994) is supported by the analytical data for the White Queen namibite. Water determination for the namibite was not possible because of the intimate intergrowth with clinobisvanite and beyerite. Eulytite is new to the State of California and probably represents a new mineral for the United States. Namibite has now been identified from about 11 different world-wide localities.

The author thanks Mr. Garth Bricker of Fallbrook, CA for providing specimens of namibite from the White Queen mine. Mr. Roland Reed of El Cajon, CA provided specimens of clinobisvanite-eulytite from the Elizabeth R mine. Mr. Blue Sheppard of Pala, CA provided specimens from the Stewart mine.

References

Foord, E.E., Spaulding, L.B., Jr., Mason, R.A. and Martin, R.F. (1989) Mineralogy and paragenesis of the Little Three mine pegmatites, Ramona district, San Diego

- County, California. *Mineral. Record*, **20**, 101–27.
- Foord, E.E., London, D., Kampf, A.R., Shigley, J.E. and Snee, L.W. (1991) Gem-bearing pegmatites of San Diego County, California. In: *Geological Excursions in southern California and Mexico* (M.J. Walawender and B.B. Hanan, eds.) Guidebook for the 1991 Annual Meeting Geological Society of America, San Diego, CA., Oct. 21–24, 1991, 128–46.
- Fron del, C. (1943) New data on agricolite, bismoclite, koechlinite, and the bismuth arsenates. *Amer. Mineral.*, **28**, 536–40.
- Jahns, R.H. (1979) Gem-bearing pegmatites in San Diego County, California: The Stewart mine, Pala district and the Himalaya mine, Mesa Grande district. In *Mesozoic Crystalline rocks*. (P.L. Abbott and V.R. Todd, eds.) San Diego, California, San Diego State University, Dept. of Geological Sciences, pp. 3–38.
- Jahns, R.H. and Wright, L.A. (1951) Gem- and lithium-bearing pegmatites of the Pala district, San Diego County, California. *Calif. Div. of Mines Spec. Report 7A*, 72 pp.
- Mrázek, Z., Veselovský, F., Hloušek, J., Moravcová, H. and Ondruš, P. (1994) Redefinition of namibite, $\text{Cu}(\text{BiO})_2\text{VO}_4\text{OH}$. *Neues Jahrb. Mineral., Mh.*, 481–8.
- Shigley, J.E., Kampf, A.R., Foord, E.E. and London, D. (1986) *Fieldtrip guidebook to gem pegmatites of southern California*, 14th General Meeting of the International Mineralogical Association, July, 1986, Stanford University, Stanford, California. 39 pp.

[Manuscript received 3 April 1995]

© Copyright the Mineralogical Society

KEYWORDS: clinobisvanite, eulytite, namibite, Pala pegmatite district, California, bismuth minerals.